

Attorney Docket No.: 05725.0441-00000

UNITED STATES PATENT APPLICATION

OF

CHRISTINE RONDEAU

FOR

COMPOSITION FOR DYEING KERATIN FIBERS WITH A  
CATIONIC DIRECT DYE AND A THICKENING POLYMER

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000

The invention relates to a composition for dyeing keratin fibers, in particular human keratin fibers such as the hair, comprising, in a medium which is suitable for dyeing, at least one cationic direct dye of given formula and at least one thickening polymer comprising at least one sugar unit.

The invention also relates to the dyeing processes and dyeing devices using the composition.

Two types of dyeing may be distinguished in the haircare sector.

The first is semi-permanent or temporary dyeing, or direct dyeing, which uses dyes capable of giving the hair a natural coloration, a more or less pronounced color change which may withstand shampooing several times. These dyes are also known as direct dyes; they can be used with or without an oxidizing agent. In the presence of an oxidizing agent, the aim is to obtain lightening dyeing. Lightening dyeing is carried out by applying a mixture, prepared at the time of use, of a direct dye and an oxidizing agent to the hair, and makes it possible in particular to obtain, by lightening the melanin in the hair, an advantageous effect such as a unified color in the case of grey hair, or to bring out the color in the case of naturally pigmented hair.

The second is permanent dyeing or oxidation dyeing. This is carried out with so-called "oxidation" dyes comprising oxidation dye precursors and couplers. Oxidation dye precursors, commonly known as "oxidation bases", are compounds which are initially colorless or weakly colored which develop their

10  
663020-5041000

15

20

US OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000

5 dyeing power on the hair in the presence of oxidizing agents added at the time of use, leading to the formation of colored compounds and dyes. The formation of these colored compounds and dyes results either from an oxidative condensation of the "oxidation bases" with themselves or from an oxidative condensation of the oxidation bases with coloration-modifying compounds commonly known as "couplers", which are generally present in the dye compositions used in oxidation dyeing.

10 It is known practice to add direct dyes to oxidation dyes in order to vary the shades obtained with the said oxidation dyes or to enrich the shades with glints.

15 Among the cationic direct dyes available in the sector of dyeing keratin fibers, in particular human keratin fibers, the compounds whose structure is developed in the text hereinbelow are already known; nevertheless, these dyes lead to colorations which have characteristics that could still be improved, such as the intensity, the homogeneity of the color distributed along the fiber, in which case the coloration is said to be too selective, and the staying power, in terms of the resistance to the various attacking factors to which the hair may be subjected (light, bad weather, shampooing).

After considerable research conducted in this matter, the inventor has now discovered that it is possible to obtain novel compositions for dyeing keratin

668020-5044100


fibers which are capable of giving more intense and yet unselective colorations which show good resistance to the various attacking factors to which the hair may be subjected, by combining at least one thickening polymer comprising at least one sugar unit with at least one known cationic direct dye of the prior art, which have the respective formulae defined below.

This discovery forms the basis of the present invention.

A first subject of the present invention is thus a composition for dyeing keratin fibers, and in particular human keratin fibers such as the hair, containing, in a medium which is suitable for dyeing, (i) at least one cationic direct dye whose structure corresponds to formulae (I) to (III') defined below, characterized in that it also contains (ii) at least one thickening polymer comprising at least one sugar unit.

(i) The cationic direct dye which can be used according to the present invention is a compound chosen from those of formulae (I), (II), (III) and (III') below:

a) the compounds of formula (I) below:



in which:

D represents a nitrogen atom or a -CH group,

R<sub>1</sub> and R<sub>2</sub>, which may be identical or different, represent a hydrogen atom; a C<sub>1</sub>-C<sub>4</sub> alkyl radical which can be substituted with a -CN, -OH or -NH<sub>2</sub> radical or form, with each other or a carbon atom of the benzene ring of formula (I), a heterocycle optionally containing oxygen or nitrogen, which can be substituted with one or more C<sub>1</sub>-C<sub>4</sub> alkyl radicals; a 4'-aminophenyl radical,

R<sub>3</sub> and R'<sub>3</sub>, which may be identical or different, represent a hydrogen atom, a halogen atom chosen from chlorine, bromine, iodine and fluorine, a cyano radical, or a C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy or acetyloxy radical,

X<sup>-</sup> represents an anion preferably chosen from chloride, methyl sulphate, perchlorate and acetate,

A represents a group chosen from the structures A1 to A19 below:
















658020 50161260

5


10

LAW OFFICES


FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000




00349405, 020899  
0668020, 00764460



A<sub>16</sub>




A<sub>17</sub>



A<sub>18</sub>

and




A<sub>19</sub>

in which R<sub>4</sub> represents a C<sub>1</sub>-C<sub>4</sub> alkyl radical which can be substituted with a hydroxyl radical and R<sub>5</sub> represents a C<sub>1</sub>-C<sub>4</sub> alkoxy radical, with the proviso that when D represents -CH, when A represents A<sub>4</sub> or A<sub>13</sub> and when R<sub>3</sub> is other than an alkoxy radical, then R<sub>1</sub> and R<sub>2</sub> do not simultaneously denote a hydrogen atom;

b) the compounds of formula (II) below:

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000



in which:

$R_6$  represents a hydrogen atom or a  $C_1$ - $C_4$  alkyl radical,

$R_7$  represents a hydrogen atom, an alkyl radical which can be substituted with a -CN radical or with an amino group, a 4'-aminophenyl radical or forms with  $R_6$  a heterocycle optionally containing oxygen and/or nitrogen, which can be substituted with a  $C_1$ - $C_4$  alkyl radical,


$R_8$  and  $R_9$ , which may be identical or different, represent a hydrogen atom, a halogen atom such as bromine, chlorine, iodine or fluorine, a  $C_1$ - $C_4$  alkyl or  $C_1$ - $C_4$  alkoxy radical or a -CN radical,

$X^-$  represents an anion preferably chosen from chloride, methyl sulphate, and




B represents a group chosen from the structures B<sub>1</sub> to B<sub>6</sub> below:






B4




B5




B6

in which  $R_{10}$  represents a  $C_1$ - $C_4$  alkyl radical,  $R_{11}$  and  $R_{12}$ , which may be identical or different, represent a hydrogen atom or a  $C_1$ - $C_4$  alkyl radical;

c) the compounds of formulae (III) and (III') below:



(III)



(III')

in which:

R<sub>13</sub> represents a hydrogen atom, a C<sub>1</sub>-C<sub>4</sub> alkoxy radical, a halogen atom such as bromine, chlorine, iodine or fluorine, or an amino radical,

R<sub>14</sub> represents a hydrogen atom, a C<sub>1</sub>-C<sub>4</sub> alkyl radical or forms, with a carbon atom of the benzene ring, a heterocycle optionally containing oxygen and/or substituted with one or more C<sub>1</sub>-C<sub>4</sub> alkyl groups,

R<sub>15</sub> represents a hydrogen atom or a halogen atom such as bromine, chlorine, iodine or fluorine,

R<sub>16</sub> and R<sub>17</sub>, which may be identical or different, represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl radical,

D<sub>1</sub> and D<sub>2</sub>, which may be identical or different, represent a nitrogen atom or a -CH group,

5  
6660/01-5076760

10

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000


m = 0 or 1,

it being understood that when R<sub>13</sub> represents an unsubstituted amino group, then

D<sub>1</sub> and D<sub>2</sub> simultaneously represent a -CH group and m = 0,


X<sup>-</sup> represents an anion preferably chosen from chloride, methyl sulphate and acetate,

E represents a group chosen from the structures E<sub>1</sub> to E<sub>8</sub> below:




E1

;




E2


;




E3




E4



E5




E6



E7


and



E8

in which R' represents a C<sub>1</sub>-C<sub>4</sub> alkyl radical;

when m = 0 and when D<sub>1</sub> represents a nitrogen atom, then E can also denote a group of structure E9 below:



in which R' represents a C<sub>1</sub>-C<sub>4</sub> alkyl radical.

In the structures (I) to (III) and (III') defined above, the C<sub>1</sub>-C<sub>4</sub> alkyl or alkoxy group preferably denotes methyl, ethyl, butyl, methoxy or ethoxy.

The cationic direct dyes of formulae (I), (II), (III) and (III') which can be used in the dye compositions in accordance with the invention are known compounds and are described, for example, in patent applications WO 95/01772, WO 95/15144 and EP-A-0,714,954, the disclosure of each of which is hereby specifically incorporated herein by reference.







Among the cationic direct dyes of formula (I) which can be used in the dye compositions in accordance with the invention, mention may be made more particularly of the compounds corresponding to the structures (I1) to (I54) below:

560020-5076260

10

LAW OFFICES






FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000



05725.0441-00000






LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000



665020-50764660















668040 50764E60

LAW OFFICES






FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000








663020-5076166








668040-50764260








063020-50761E60








05725.0441-00000






666020-50764E60





and






05725.0441-00000






LAW OFFICES

FINNEGAN, HENDERSON,  
 FARABOW, GARRETT,  
 & DUNNER, L.L.P.  
 1300 I STREET, N. W.  
 WASHINGTON, D. C. 20005  
 202-408-4000



Among the compounds of structures (I1) to (I54) described above, the ones most particularly preferred are the compounds corresponding to the structures (I1), (I2), (I14) and (I31).

Among the cationic direct dyes of formula (II) which can be used in the dye compositions in accordance with the invention, mention may be made more particularly of the compounds corresponding to the structures (II1) to (II9) below:






669020" 5076hE60



and




000020" 50764260

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000

Among the cationic direct dyes of formula (III) which can be used in the dye compositions in accordance with the invention, mention may be made more particularly of the compounds corresponding to the structures (III1) to (III18) below:









;





668020-50764E60

LAW OFFICES


FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000







; and



05725.0441-00000




LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000



Among the specific compounds of structures (III1) to (III18) described above, the ones most particularly preferred are the compounds corresponding to the structures (III4), (III5) and (III13).

Among the cationic direct dyes of formula (III') which can be used in the dye compositions in accordance with the invention, mention may be made more particularly of the compounds corresponding to the structures (III'1) to (III'3) below:



668020-50764E60

000020" 50764E60

The cationic direct dye(s) used according to the invention preferably represent(s) from 0.001 to 10% by weight approximately relative to the total weight of the dye composition and even more preferably from 0.005 to 5% by weight approximately relative to this weight.

(ii) The thickening polymer comprising at least one sugar unit which can be used according to the present invention is chosen from:

(ii)<sub>1</sub> - nonionic guar gums;

(ii)<sub>2</sub> - biopolysaccharide gums of microbial origin, such as scleroglucan or xanthan gums;

(ii)<sub>3</sub> - gums derived from plant exudates, such as gum arabic, ghatti gum, karaya gum, gum tragacanth, carrageenan gum, agar gum and carob gum;

(ii)<sub>4</sub> - pectins;

(ii)<sub>5</sub> - alginates;

(ii)<sub>6</sub> - starches; and

(ii)<sub>7</sub> - hydroxyalkylcelluloses and carboxyalkylcelluloses.

For the purposes of the present invention, the expression "sugar unit" denotes a monosaccharide portion (i.e., monosaccharide or oside or simple sugar) or an oligosaccharide portion (short chains formed from the linking of monosaccharide units, which may be different) or a polysaccharide portion (long chains consisting of monosaccharide units, which may be different, i.e., polyholosides or polyosides (homopolyosides or heteropolyosides). The saccharide units can also be substituted with alkyl, hydroxyalkyl, alkoxy, acyloxy or carboxyl groups.

The nonionic guar gums can be modified or unmodified. The unmodified guar gums are, for example, the products sold under the name Vidogum GH 175 by the company Unipectine and under the name Jaguar C by the company

5

10

15

20

Meyhall.

According to the present invention, it is preferred to use nonionic guar gums modified with C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl groups.

Among the hydroxyalkyl groups which may be mentioned, for example, are hydroxymethyl, hydroxyethyl, hydroxypropyl and hydroxybutyl groups. These guar gums are well known in the state of the art and can be prepared, for example, by reacting the corresponding alkene oxides such as, for example, propylene oxides, with the guar gum so as to obtain a guar gum modified with hydroxypropyl groups.

The degree of hydroxyalkylation, which corresponds to the number of alkylene oxide molecules consumed by the number of free hydroxyl functions present on the guar gum, preferably ranges from 0.4 to 1.2.

Such nonionic guar gums optionally modified with hydroxyalkyl groups are sold, for example, under the trade names Jaguar HP8, Jaguap HP60 and Jaguar HP120, Jaguar DC 293 and Jaguar HP 105 by the company Rhône-Poulenc (Meyhall) or under the name Galactasol 4H4FD2 by the company Aqualon.

The biopolysaccharide gums of microbial origin, such as the scleroglucan or xanthan gums, the gums derived from plant exudates such as gum arabic, ghatti gum, karaya gum, gum tragacanth, carrageenan gum, agar gum and carob gum, the hydroxyalkylcelluloses and carboxymethylcelluloses, pectins,

alginates and starches are well known to those skilled in the art and are described in particular in the book by Robert L. Davidson entitled "Handbook of Water soluble gums and resins" published by McGraw Hill Book Company (1980), the disclosure of which is specifically incorporated by reference.

5           Among these gums, the scleroglucans more particularly used according to the present invention are represented by the products sold under the name Actigum CS by the company Sanofi Bio Industries and in particular Actigum CS 11, and under the name Amigel by the company Alban Muller International. Other scleroglucans, such as the one treated with glyoxal in French patent application No. 2,633,940, can also be used, the disclosure of which is incorporated by reference.

10           The xanthan gums more particularly used according to the present invention are represented by the products sold under the names Keltrol, Keltrol T, Keltrol TF, Keltrol BT, Keltrol RD and Keltrol CG by the company Nutrasweet Kelco, or under the names Rhodicare S and Rhodicare H by the company Rhodia Chimie.

15           The hydroxyalkylcelluloses are more particularly hydroxyethylcelluloses, such as those sold under the names Cellosize QP3L, Cellosize QP4400 H, Cellosize QP30000H, Cellosize HEC30000A and Cellosize Polymer PCG10 by the company Amerchol, or Natrosol 250HHR, Natrosol 250 MR, Natrosol 250M,

Natrosol 250HHXR, Natrosol 250HHX, Natrosol 250HR and Natrosol HX by the company Hercules, or Tylose H1000 by the company Hoechst.

The hydroxyalkylcelluloses are also, more particularly, hydroxypropylcelluloses such as the products sold under the names Klucel EF, Klucel H, Klucel LHF, Klucel MF and Klucel G by the company Aqualon.

Among the carboxyalkylcelluloses preferably used is carboxymethylcellulose, for which mention may be made of the products sold under the names Blanose 7M8/SF, Blanose Raffinée 7M, Blanose 7LF, Blanose 7MF, Blanose 9M31F, Blanose 12M31XP, Blanose 12M31P, Blanose 9M31XF, Blanose 7H, Blanose 7M31 and Blanose 7H3SXF by the company Aqualon, or Aquasorb A500 and Ambergum 1221 by the company Hercules, or Cellogen HP810A and Cellogen HP6HS9 by the company Montello, or Primellose by the company Avebe.

The thickening polymers (ii) used in the compositions of the present invention are preferably present in a proportion of from 0.01 to 10% by weight approximately, in particular in a proportion of from 0.1 to 5% by weight approximately, relative to the total weight of the dye composition applied to the keratin fibers.

The medium which is suitable for dyeing (or support) generally comprises water or a mixture of water and at least one organic solvent to dissolve the


compounds which would not be sufficiently water-soluble. As organic solvents, mention may be made, for example, of C<sub>1</sub>-C<sub>4</sub> lower alkanols such as ethanol and isopropanol; aromatic alcohols such as benzyl alcohol, as well as similar products and mixtures thereof.

5           The solvents can be present in proportions preferably from 1 to 40% by weight approximately relative to the total weight of the dye composition, and even more preferably from 5 to 30% by weight approximately.

10           The pH of the dye composition in accordance with the invention is generally approximately from 2 to 11 and preferably approximately from 5 to 10. It can be adjusted to the desired value using acidifying or basifying agents usually used for dyeing keratin fibers.

15           Among the acidifying agents, mention may be made, by way of example, of inorganic or organic acids such as hydrochloric acid, orthophosphoric acid, sulphuric acid, carboxylic acids such as acetic acid, tartaric acid, citric acid and lactic acid, and sulphonic acids.

          Among the basifying agents, mention may be made, by way of example, of aqueous ammonia, alkaline carbonates, alkanolamines such as mono-, di- and triethanolamine and derivatives thereof, sodium hydroxide, potassium hydroxide and the compounds of formula (IV) below:



in which W is a propylene residue optionally substituted with a hydroxyl group or a C<sub>1</sub>-C<sub>6</sub> alkyl radical; R<sub>18</sub>, R<sub>19</sub>, R<sub>20</sub> and R<sub>21</sub>, which may be identical or different, represent a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub> alkyl or C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl radical.

In addition to the cationic direct dye(s) (i) defined above, the dye composition in accordance with the invention can comprise one or more additional direct dyes which can be chosen, for example, from nitrobenzene dyes, anthraquinone dyes, naphthoquinone dyes, triarylmethane dyes, xanthene dyes and azo dyes which are non-cationic.

When it is intended for oxidation dyeing, the dye composition in accordance with the invention comprises, in addition to the cationic direct dye(s) (i), one or more oxidation bases chosen from the oxidation bases conventionally used for oxidation dyeing and among which mention may be made in particular of para-phenylenediamines, bis(phenyl)alkylenediamines, para-aminophenols, ortho-aminophenols and heterocyclic bases.

When they are used, the oxidation base(s) preferably represent(s) from 0.0005 to 12% by weight approximately relative to the total weight of the dye composition, and even more preferably from 0.005 to 6% by weight



approximately relative to this weight.

When it is intended for oxidation dyeing, the dye composition in accordance with the invention can also comprise, in addition to the cationic direct dye (i) and the thickening polymer (ii) as well as the oxidation bases, one or more couplers so as to modify the shades obtained or to enrich them with glints, by using the cationic direct dye(s) (i) and the oxidation base(s).

The couplers which can be used in the dye composition in accordance with the invention can be chosen from the couplers used conventionally in oxidation dyeing and among which mention may be made in particular of meta-phenylenediamines, meta-aminophenols, meta-diphenols and heterocyclic couplers.

When it is (they are) present, the coupler(s) preferably represent(s) from 0.0001 to 10% by weight approximately relative to the total weight of the dye composition, and even more preferably from 0.005 to 5% by weight approximately relative to this weight.

The dye composition in accordance with the invention can also contain various adjuvants conventionally used in compositions for dyeing the hair, such as antioxidants, penetrating agents, sequestering agents, fragrances, buffers, dispersing agents, surfactants, film-forming agents, ceramides, preserving agents, screening agents, such as sunscreens, and opacifiers.

Needless to say, a person skilled in the art will take care to select this (these) optional complementary compound(s) such that the advantageous properties intrinsically associated with the dye composition in accordance with the invention are not, or are not substantially, adversely affected by the addition(s) envisaged.

The dye composition according to the invention can be in various forms, such as in the form of liquids, shampoos, creams or gels or any other form which is suitable for dyeing keratin fibers, and in particular human hair. It can be obtained by mixing, at the time of use, a composition, which may be pulverulent, containing the cationic direct dye(s) with a composition containing the thickening polymer (ii) according to the invention.

When the combination of the cationic direct dye (i) and the thickening polymer (ii) according to the invention is used in a composition intended for oxidation dyeing (in which case one or more oxidation bases are used, optionally in the presence of one or more couplers) or when it is used in a composition intended for lightening direct dyeing, then the dye composition in accordance with the invention also comprises at least one oxidizing agent chosen, for example, from hydrogen peroxide, urea peroxide, alkali metal bromates, persalts such as perborates and persulphates, and enzymes such as peroxidases, lactases and two-electron oxidoreductases. It is particularly preferred to use

hydrogen peroxide or enzymes.

Another subject of the invention is a process for dyeing keratin fibers, and in particular human keratin fibers such as the hair, using the dye composition as defined above.

5 According to a first variant of this dyeing process in accordance with the invention, at least one dye composition as defined above is applied to the fibers, for a period which is sufficient to develop the desired coloration, after which the fibers are rinsed, optionally washed with shampoo, rinsed again and dried.

10 The time required to develop the coloration on the keratin fibers is generally from 3 to 60 minutes and even more specifically from 5 to 40 minutes.

According to a second variant of this dyeing process in accordance with the invention, at least one dye composition as defined above is applied to the fibers, for a period which is sufficient to develop the desired coloration, without final rinsing.

15 According to one specific embodiment of this dyeing process, and when the dye composition in accordance with the invention comprises at least one oxidation base and at least one oxidizing agent, the dyeing process comprises a first step which comprises separately storing, on the one hand, a composition (A1) comprising, in a medium which is suitable for dyeing, at least one cationic direct dye (i) as defined above and at least one oxidation base, and, on the other

hand, a composition (B1) comprising, in a medium which is suitable for dyeing, at least one oxidizing agent, and then in mixing them together at the time of use, after which this mixture is applied to the keratin fibers, the composition (A1) or the composition (B1) containing the thickening polymer (ii) as defined above.

5

According to another specific embodiment of this dyeing process, and when the dye composition in accordance with the invention comprises at least one oxidizing agent, the dyeing process comprises a first step which consists in separately storing, on the one hand, a composition (A2) comprising, in a medium which is suitable for dyeing, at least one cationic direct dye (i) as defined above, and, on the other hand, a composition (B2) comprising, in a medium which is suitable for dyeing, at least one oxidizing agent, and then in mixing them together at the time of use, after which this mixture is applied to the keratin fibers, the composition (A2) or the composition (B2) containing the thickening polymer as defined above.

15

Another subject of the invention is a multi-compartment dyeing device or dyeing "kit" or any other multi-compartment packaging system, a first compartment of which comprises the composition (A1) or (A2) as defined above and a second compartment of which comprises the composition (B1) or (B2) as defined above. These devices can be equipped with means for dispensing the desired mixture onto the hair, such as the devices described in patent FR

20

5

## EXAMPLES

The four direct dyeing compositions given in the table below were prepared:

WATERBURY

	Example 1	Example 2	Example 3	Example 4
Cationic direct dye of formula (I1)	0.2			
Cationic direct dye of formula (I2)		0.2		
Cationic direct dye of formula (I14)			0.2	
Cationic direct dye of formula (I31)				0.15
Hydroxyethylcellulose sold under the name Natrosol 250 HHR by the company Aqualon	1.0 AM*			
Carboxymethylcellulose sold under the name Blanose 7M by the company Aqualon		1.0 AM*		
Guar gum sold under the name Vidogum GH175 by the company Unipeptine			1.0 AM*	
Scleroglucan gum sold under the name Amigel by the company Alban Muller International				1.0 AM*
Ethanol	10	10	10	10
2-Amino-2-methyl-1-propanol qs	pH 9	pH 9	pH 9	pH 9
Demineralized water qs	100	100	100	100

5

10

15

20

25

660020 50767260

LAW OFFICES

FINNEGAN, HENDERSON,  
FARABOW, GARRETT,  
& DUNNER, L.L.P.  
1300 I STREET, N. W.  
WASHINGTON, D. C. 20005  
202-408-4000

AM\* denotes active material

AM\* denotes active material

The above compositions were each applied for 30 minutes to locks of natural grey hair containing 90% white hairs. The locks of hair were then rinsed, washed with a standard shampoo and then dried.

The locks were dyed in the following shades:

Examples	Shades obtained
1	Bright red
2	Bright red
3	Bright orange
4	Bright purple

10 668020 50764E60